

Antifungal Activity of Titanium Dioxide Photocatalysis Against *Fusarium oxysporum* f.sp.*lycopersici* .

*Rajaa A. AL-Anbagi **Fakhir E. Hameed *Fatima Al-Zahraa G.Gassim

*College of Science for Women

**College of Agriculture

Univ. Babylon,Iraq

ABSTRACT

Studies were carry out to detect the efficacy of titanium dioxide (TiO₂) photocatalysis combined with light (mercury lamp ,160 W) on number of colony forming unites (CFUs) and dry weight of biomass of fungus of *Fusarium oxysporum* f.sp. *lycopersici* Schlecht in photocatalytic reaction cell during different exposure periods of light . The results showed that TiO₂ combined with light caused significantly reduced CFUs to 65.66 ,12.66, 4 and 4.66 CFUs/ 0.5 ml after periods of time 30,60,90 and 120 min respectively compared with control in the dark without TiO₂ ,while TiO₂ alone didn't effect on CFUs compared with control in the dark . When light was present along time with TiO₂ ,it was found the survival ratio reduction into 1.93 and 2.2 % after 90 and 120 min., while rate of photo killing of TiO₂ (slope) was 1.5531 CFUs /min . Also observed TiO₂ combined with light was reduced significantly in dry weight of biomass of *F.oxysporum* f.sp.*lycopersici* to 42 and 50 mg /30ml after exposure it into periods 60 and 120 min respectively compared with control in the dark.

INTRODUCTION

The element titans (Titanium) was discovered in 1791 by William Gregor, in England . Martin Klaproth, Later named it titanium and he was only able to produce titanium dioxide . In nature its never occurrence pure .It found with contaminant metal such as iron (Higgin ,1973). Titanium dioxide (TiO₂) is a white powder ,occurs in three crystalline forms ,anatase ,rutile and brookite . Boiling point 2972 C° in soluble , molecular weight 79.87 g /mol , density 4.23 g /cm³ (Fox and Dulay,1993). It have important properties is photocatalysts when UV illuminated it with wave length less than 385 nm . Photocatalysts generate a strong oxidizing power and could be decompose organic and inorganic compounds by oxidation or reduction (Higgin ,1973; Lee, 2004) . The two crystalline forms of titanium dioxide , anatase and rutile have property photocatalysis the least it has been found to most active form (Higgin ,1973) .

Titanium dioxide(TiO₂) is a multifaceted compound , its the stuff that makes tooth paste white and paint opaque because non-toxic for human therefore its used in cosmetics products and in special pharmaceuticals (Doll and Frimmel, 2005). Also titanium dioxide has been widely utilized as self – cleaning , self sterilizing material for coating clinical tools , items for use in hospital (Fujishima *et al.*,1999) and in the purification of water and air on surfaces from microorganisms such as bacteria,viruses,protozoa and fungi (Lee, 2004; Lonnen *et al.* ,2005).

In 1985 the first research work on the microbiocidal effect of titanium dioxide on microbial cell of *Escherichia coli* was found in water and it could be killed by contact with a TiO₂-pt catalyst upon illumination with near – UV light for 60 to 120 min (Matsunaga *et al.* ,1985).Since then sub sequently has been intensively conducted on a wide spectrum of organisms primarily with bacteria and tumor cells (Blake *et al.*,1999 ; Lonnen *et al.* ,2005).

Saito *et al.* (1992) and Maness *et al.* (1999) have explained , that particles come into contact with the gram positive bacteria as *Micrococcus luteus* and *Streptococcus sorbinus* , when irradiation titanium dioxide .The microbial surface was the primary target of initial oxidative

attack reactive oxygen species (ROS) such as hydroxyl radical (OH^\cdot), superoxide (O_2^\cdot) and hydrogen peroxide (H_2O_2) were generated on the irradiated titanium dioxide surface. Susceptibility four kinds of organisms (*E. coli*, *Lactobacillus acidophilus*, *Saccharomyces cerevisiae* and *Chlorella vulgaris*) to killing by the photocatalytic effect was observed when it was compared with using platinized titanium dioxide and a metal halide lamp. The photocatalytic method has found its application also in the degradation of toxins secreted to water by bacteria and unicellular protozoa and degradation of algae, bacteria, viruses and protozoa which normally found in it (Matsunaga *et al.*, 1988; Robertson *et al.*, 1997; Lawton *et al.*, 1999; Makowski and Wardas, 2001). Also its application to sterilize selected food borne pathogenic bacteria such as *Salmonella choleraesuis*, *Vibrio parahaemolyticus*, *Listeria monocytogenes* and *Pseudomonas sp* (Kim *et al.*, 2006).

The fungicidal effect of the TiO_2 photocatalytic ozonation reaction for control of *Diaporthe actinidiae* on kiwifruit and it was used to control post harvest storage rots in kiwifruit and decompose residual the fungicides (Hur *et al.*, 2005). The antifungal activity of TiO_2 photocatalytic reaction in the form of TiO_2 powder and TiO_2 coated on plastic film against *Penicillium expansum* (air borne fungus) in vitro and in apple fruit was recorded (Maneerat and Hayata, 2006). When stimulated solar and solar photocatalytic exposure 870 W/m^2 in the 300 nm -10 μm range/ 200 W/m^2 in the 300-400 nm UV range, its reduced in viability against trophozoite stage of protozoa of *Acanthamoeba polyphaga*, *Candida albicans*, *F. solani* and *Bacillus* found in water (Lonnen *et al.*, 2005). Also was obtained ability of solar only and solar photocatalytic (TiO_2) of five wild strain of *Fusarium* which was successfully achieved (Sichel *et al.*, 2007).

In this study, we investigated the effect of TiO_2 powder and TiO_2 photocatalysis on the fungus of *F. oxysporum* that as plant pathogen and producer to mycotoxin (fumonisin) that is cytotoxic effect to several mammalian cell lines (Abbas *et al.*, 1998; Dignani and Anaissie, 2004). This a study a first in Iraq about TiO_2 photocatalysis in fungi.

MATERIAL AND METHODS

Isolation and preparation of the fungus

Isolates of *Fusarium oxysporum* f.sp. *lycopersici* Schlecht was isolated from infected stem of tomato *Lycopersicon esculentum* by cultured some sections of infected parts after its surface sterile on Petri dish contain 20 ml potato dextrose agar media (PDA), the fungus was purified and identification by protocol Hansen and Smith (1932). Spores suspension were prepared from the fungus by suspended spores of *F. oxysporum* f.sp. *lycopersici* with sterile distilled water and it was counted this spores per ml by a hemocytometer (Kerley and Solymosy, 1974) for using in following experiments.

Titanium dioxide (TiO_2)

The photocatalyst titanium dioxide powder was supplied by Degussa company P-25 (Japan) particles with an average composition of 75% anatase and 25% rutile. Physical properties of TiO_2 crystallite were characterized by BET (Brunauer – Emmett – Teller) analysis, which is non – porous, with a surface area about of $55 \text{ m}^2/\text{g}$. It has a particle size of 0.03 micron and an average particles diameter of 21 nm (Gassim *et al.*, 2004; Coleman *et al.*, 2005). The P-25 titanium dioxide Degussa has become the standard for photo reactivity in water, air purification and bactericidal (Blake *et al.*, 1999; Maness *et al.*, 1999; Makowski and Wardas, 2001). This compound was used for all experiments and stored at room temperature.

Photocatalytic reaction cells

The photocatalytic reactor consisted of low pressure mercury lamp type Emkay (160 W) wave length between 360-750 nm was used as source of irradiation, photo cell contain the

vessel (35ml)with quartz window (2cm²)as reaction vessel ,oxygen pump .The light lamp was centered to illuminated properly the inner of vessel and the temperature was controlled at 25C° by using thermo-circulator (Desaga Frigostat) during the photocatalytic reaction .




Fig. 1. Schematic diagram of the experimental apparatus for photocatalytic reaction .(A) gas container, (B) gas flowmeter,(C) circulating water thermostat ,(D) magnetic stirrer, (E) quartz photo cell, (F) windows quartz, (H) low pressure mercury lamp, (I) power supply unit .

Effect photocatalytic reaction of TiO₂ on numbers of colony forming unites (CFUs) of *F.oxysporum* f.sp. *lycopersici*

To investigate the antifungal activity of TiO₂ Photocatalysis, 120 ml spores suspension have 8×10^2 spore /ml were freshly prepared with sterile distilled water for all treatments, aliquots of 30 ml from spores suspension per each treatment, four treatments were carried out (i) the control treatment in the dark by adding 30 ml of spores suspension to the photocatalytic reaction cell , quartz window were covered with black cover without TiO₂ . (ii) treatment in the light , , spores suspension was prepared by the same method for the first treatments but photocatalytic reaction cell were exposed to the light without TiO₂ . (iii) the TiO₂ alone treatment was prepared with 10 mg of TiO₂ adding into 30 ml spores suspension, quartz window were covered with black cover . (iv) the last treatment 10 mg of TiO₂ adding into 30 ml of spores suspension and its added into the photocatalytic reaction cell then its exposure to the light (Maneerat and Hayata, 2006). Spores suspensions of all treatments were stirring by using magnetic stirrer. Oxygen gas was passed with rate 10 cm³ / min to the photocatalytic reaction cell .Temperature was controlled in 25 C° using the thermo – circular during the photocatalytic reaction cell in all treatments .The treatments samples were collected from the reaction cell every subsequent 30 minute .For each sampling,2 ml of the suspension was draws by using a syringe with along pliable needle from the reaction cell for all treatments after 0,30,60 ,90 and 120 min ,then the treatments samples were centrifuged at 1000 rpm for 5 min to separate the solid catalyst,0.5 ml of supernatant immediately added into Petri dishes (9cm diameter) , than 20 ml of PDA media poured into Petri dish with trireplicates per each treatments .

The Petri dishes were incubated in the dark at 30 C° \pm 2 for 48 hours. The numbers of colony forming unites of *F. oxysporum* per each plate were counted (Leonard and Blackford

,1949 ; Ohmori and Gottlieb ,1965) .The survival ratio (%) of *F.oxysporum* f.sp. *lycopersici* in aqueous solution and rate of photokilling of TiO₂ (slope) were calculated .

Effect photocatalytic reaction of TiO₂ on dry weight of *F.oxysporum* f.sp. *lycopersici*

The experiment was carried out with four treatments as well as previous experiment (control in the dark, light ,TiO₂ alone, and TiO₂ with light).Spores suspensions of all treatments were stirring by using magnetic stirrer.Oxygen gas was passed with rate 10 cm³ /min to the photocatalytic reaction cell .Temperature was controlled in 25 C° using the thermo – circular during the photocatalytic reaction cell in all treatments, 2 ml of the suspension were draws by using a syringe with along pliable needle from the reaction cell for all treatments after 60 and 120 min, then the treatmentse samples were centrifuged at 1000 rpm for 5 min to separate the solid catalyst,0.5 ml of supernatant immediately added into bottle have 30 ml PDB(potato dextrose broth) with trireplicates per each treatment ,the treatments were incubated in the dark at 30C° in 2 for 14 days. Dry weights of biomass of the fungus were obtained by filtrated cultured media then drying biomass at 70 C° for 24 h (Singh *et al* .,1980).

Statistical analysis

All experiments were designed complete randomized design and data analyzed by using least squares analysis of variance (ANOVA), least significant difference (L. S. D.) test was used at the 1% and 5% level of significance (Steel and Torrie,1960).

RESULTS

Results effect of treatments (control in the dark , the light treatment ,TiO₂ alone and TiO₂ combined with light)on number of colonies forming unites(CFUs) of *Fusarium oxysporum* f.sp. *lycopersici* were showed that TiO₂ combined with light caused significantly reduced the colonies forming unites(CFUs) to 65.66, 12.66 ,4 and 4.66 CFUs /0.5 ml after exposure periods of light 30, 60, 90 and 120 min respectively compared with dark treatment .Also observed reducing in the numbers of colony forming unites in the light treatments compared with dark treatment. While TiO₂ alone didn't effect on numbers of colonies forming unites compared with control in the dark or light treatments (Table1 and Fig. 2).

Table1. Effect of the TiO₂ photocatalytic reaction on numbers of colony forming unites(CFUS) of *Fusarium oxysporum* f.sp.*lycopersici*.

Time(min)	Numbers of colony forming unites / 0.5 ml *			
	Treatments of TiO ₂			
	Control in the dark	Light treatment	TiO ₂ alone	TiO ₂ with light
30	205.00	160.66	210.66	65.66
60	213.00	141.66	201.66	12,66
90	207,00	166.33	197.33	4.00
120	202.30	174.00	217.00	4.66
Means	206.80	160.66	206.66	21.75

L.S.D. 0. 01 for treatments of TiO₂ = 24.4

L.S.D. 0. 05 for Interaction between treatment of TiO₂ and time = 33.62

* each number is mean of trireplicates.




Fig. 2. Effect of TiO₂ and light on numbers of colony forming unites(CFUs) of *Fusarium oxysporum* f.sp. *lycopercici* . (A) control in the dark after 30 min , (B) control in the dark after 60 min., (C) control in the dark after 90 min. , (D) control in the dark after 120 min., (E) light exposure after 30 min., (F) light exposure after 60 min., (G) light exposure after 90 min., (H) light exposure after 120 min , (I) TiO₂ alone in the dark after 30 min., (J) TiO₂ alone in the dark after 60 min., (K) TiO₂ alone in the dark after 90 min., (L) TiO₂ alone in the dark after 120 min. ,(M) TiO₂ combined with light exposure after 30 min., (N) TiO₂ combined with light exposure after 60 min., (O) TiO₂ combined with light exposure after 90 min., (P) TiO₂ combined with light exposure after 120 min.

Also TiO_2 photocatalytic causes reduced survival ratio of *F.oxysporum* f.sp. *lycopersici* to 31.7, 6.12, 1.93 and 2.2 % after exposure periods of light 30, 60, 90 and 120 min respectively (Fig.3). While ,rate of photo killing of TiO_2 (slope) was 1.5531 CFU /min (Fig.4).




Fig. 3. Effect of the TiO_2 photocatalyst on the survival ratio (%) of *F.oxysporum* f.sp. *lycopersici* in aqueous solution at 25 C°.




Fig . 4. The relationship between colony forming units (CFU) / 0.5 ml with time of irradiation of TiO_2 at 25 C°.

Results of dry weight of biomass of *F.oxysporum* f.sp. *lycopersici*. showed that TiO_2 combined with light caused significantly reduced of biomass to 42 and 50 mg / 30 ml after exposure period 60 and 120 min respectively compared with control treatment in the dark were 168 and 168.3 mg / 30 ml and light treatment were 156.3 and 143.3 mg /30 ml .While didn't found any significant difference between exposure periods of time and treatments of TiO_2 (Table 2) .

Table 2. Effect of TiO₂ photocatalytic reaction on dry weight of biomass of *Fusarium oxysporum* f.sp *lycopersici*.

Time(min)	Dry weight of biomass (mg)/ 30 ml *				
	Control in the dark	Light treatment	TiO ₂ alone	TiO ₂ with light	means
60	168.0	156.3	143.0	42.0	127.32
120	168.3	143.3	125.0	50.0	121.65
Means	168.1	149.8	134.0	46.0	124.47

L.S.D. 0. 01 for treatments of TiO₂ = 48

*** each number is mean of trireplicates**

Discussion

Our results of illuminated TiO₂ photocatalyst effect conformed previous researches showed that killing property of illuminated TiO₂ on other microorganisms such as *E. coli*, *Streptococci*, *Lactobacillus*, *Salmonella*, *Candida albicans*, *Saccharomyces cerevisiae* were observed (Matsunaga *et al.*,1985,1988; Saito *et al.*,1999; Maness *et al.*, 1999). Also that killing property was found to positively correlate with time, type of source was used irradiation, type of TiO₂ and organisms.

Significant effect in light treatment compared with control treatment in the dark may be attributed to inhibited effect of light on fungal growth (Grow and Gadd,1995) while TiO₂ alone didn't caused any effect on colonies forming unites and dry weight because its non toxic effect on human therefore, its using in cosmetic products and pharmaceuticals (Blake *et al.*,1999; Doll and Frimmel, 2005). The significant reduction of TiO₂ combined with light treatments of biomass may be due to low colonies from unites led to reduce biomass.

The mechanism of photokilling is,when photocatalyst titanium dioxide (TiO₂) two crystalline forms of TiO₂ have photocatalytic activity, anatase and rutile. Anatase has a forbidden band gap 3.2 eV and rutile 3.0 eV. Anatase has been found to be the most active form. The action spectrum for anatase shows a strong reduction of activity in wavelengths higher than 385 nm. The photocatalytic process includes chemical steps that produce reactive species in principal can cause fatal damage to structure or functions of microorganisms cells (Fox and Dually, 1993). So the photocatalytic TiO₂ in aqueous solution it was absorbed Ultraviolet radiation from sunlight or illuminated light source (fluorescent lamps), it will produce pairs of electrons and holes. The electron of the valence band of titanium dioxide becomes excited when illuminated by light. The excess energy of this excited electron promoted the electron to the conduction band of titanium dioxide therefore creating the negative – electron (e⁻) and positive – hole (h⁺) pair. This stage is referred as the semiconductor's (photo-excitation) state. The energy difference between the valence band and the conduction band is known as the Band G' (Fig.5) (Wong *et al.*,2006; Hoffmann *et al.*,1995).





Fig. 5. Photocatalysis mechanism of titanium dioxide (Wong *et al.*,2006)

The positive – hole of titanium dioxide breaks the apart of water molecule to form hydrogen gas(H₂) and hydroxyl radicals (OH[·]) . The negative – electron reacts with atmospheric oxygen molecule(adsorbs on the surface of TiO₂ particles) to form super oxide ions .These hydroxyl radicals contact with each other to produce hydroxyl peroxide(H₂O₂) this cycle continues when light is available of the photocatalytic system , there can also be direct photochemistry as there would be from any UV source . Mechanism of a photocatalytic process on irradiated titanium dioxide(Barbeni *et al.* , 1987) : Electron –Hole Pair Formation.



(conduction band electron and valence band hole)

Electron removal from the conduction band



Nonproductive radical reactions $O_2^{\cdot-}$



For a cell or virus in contact with the titanium dioxide surface these also be direct electron or hole transfer to the organism or one of its components . If titanium dioxide particles are small size ,they may penetrate into the cell and these processes could in the interior .Since light is an essential component of the photocatalytic system (Kamat , 1993 ; Blake *et al* ., 1999) . Hydroxyl radicals are highly reactive and therefore short – lived. Superoxide ion are more long-lived; however ,due to the negative charge they cannot penetrate the cell membrane .Upon their production on the TiO_2 surface ,both hydroxyl radicals and super oxide would have to interact immediately with the outer surface of an organism unless the TiO_2 particle has penetrate into the cell (Neiland ,1982 ;Blanco - Gálvez *et al* .,2007) (Fig.6)

Compared to hydroxyl radical and super oxide ions, hydrogen peroxide is less detriment . However, the important part for killing hydrogen peroxide can enter the cell and be activated by ferrous ion via the Fenton reaction :





Figure 6 . Schematic illustration of solar photocatalytic process for bacteria inactivation in the presence of an aqueous suspension of TiO_2 (relative size of each element is schematically represented at the bottom) (Blanco, Malato, Fernández-Ibáñez, 2007).

The photokilling of illuminated TiO_2 because the reactive oxygen species (ROS), such as $\cdot OH$, O_2 and H_2O_2 generated on the irradiated TiO_2 surface have been proposed to attach with polyunsaturated phospholipids in cell membrane of *E.coli* . Iron levels on the cell surface , in the periplasmic space or inside the cell , either as iron clusters or in iron storage proteins (such as ferritin) are significant and can serve as a source of ferrous ion . Therefore , while the TiO_2 is

being illuminated to produce H_2O_2 the Fenton reaction may take place in vivo and produce the more damaging hydroxyl radicals (Neiland, 1982; Cai *et al.*, 1991; Maness *et al.*, 1999).

When the light is turned off, any residual hydrogen peroxide would continue to interact with the iron species and generated additional hydroxyl radicals through the Fenton reaction. When both H_2O_2 and O_2^- are present, the iron-catalyzed Haber-Weiss reaction can provide a second pathway to form additional hydroxyl radicals (Youngman, 1984).



Therefore the lipid peroxidation reaction that causes a break down of the cell membrane structure and its associated functions is the mechanism underlying cell death. Because all life forms have cell membrane. Thus, the proposed killing mechanism is applicable to all cell type (Saito *et al.*, 1992; Maness *et al.*, 1999). Apart from the cell wall, there exists another possible cause of death, this one is the destructive effect of oxidative photocatalysis on RNA and DNA molecules, mainly due to hydroxyl radicals (Tachikawa *et al.*, 2008).

The antifungal activity of TiO_2 photocatalytic reaction in the form of TiO_2 powder and TiO_2 coated on plastic film against *Penicillium expansum* (air born fungus) and *Diaporthe actinidiae* a major fungal pathogen of kiwifruit (Hur *et al.*, 2005; Maneerat and Hayata, 2006). The mechanisms for antifungal effect presented by Matsunaga *et al.* 1988 they were evidenced for the oxidation of coenzyme A (COA) in *S.cereviace*, a yeast when exposed to light and platinized TiO_2 for 120 minutes under a metal halide lamp, more than 97% of intracellular COA content was lost and 42% of respiratory activity was decreased led to cell death and observed the cell membrane would have to be oxidized first under go its semi permeability. Same authors failed to detect and destruction of cell wall by photoactivated semiconductor powder but Hammel *et al.* (2002) noted the degradation of polysaccharides by OH^\cdot has also recently by them that the OH^\cdot abstracts hydrogen atoms from sugar subunits of polysaccharides, resulting in cleavage of the polysaccharide chain. We suggest for its effect on spores suspension of *F.oxysporum* may be the reactive (ROS) lead to breakdown the contains such as protein, lipid and polysaccharide (cellulose) for thin cell wall of spores first and degradation of cell membrane then effected its on other chemical compounds activates such as respiratory or, and TiO_2 particles may be affection on structures of DNA and RNA for *F.oxysporum* spores all, or some factors led to photokilling it and prevent the germination of spores therefore didn't formation of fungal colonies.

REFERENCE

- Abbas, HK; Shier, WT; Seo, J-A and Lee, Y.W. 1998. Phytotoxicity and cytotoxicity of fumonosin C and P series of mycotoxins from *Fusarium* spp fungi. *Toxicon*. 36: 2033-2037.
- Barbeni, M.; Morello, M. and Pelizzetti, E. 1987. Sunlight photodegradation of 2,4,5-Trichlorophenoxy acetic acid and 2,4,5-trichlorophenol on TiO_2 . Pergamon J. Ltd. 16 (6): 1165-1179.
- Blake, D.; Maness, P-C.; Huang, Z.; Wolfrum, E.; Huang, J. and Jacoby, W. 1999. Application of the photocatalytic chemistry of titanium dioxide to disinfection and the killing of cancer cells. *Separation and purification methods*. 28(1): 1-50.
- Blanco-Gálvez, J.; Fernández-Ibáñez, P.; and Malato-Rodríguez, S. 2007. Solar photocatalytic detoxification and disinfection of water: recent overview. *Solar Energy Engineering*. 129 (1): 4-15.
- Cai, R.; Hashimoto, K.; Kubota, Y.; and Fujishima, A. 1991. Photokilling of malignant cells with ultra fine TiO_2 powder. *Bull. Chem. Soc. Jpn.* 64: 1268-1273.

- Coleman, H.M.; Marquis ,C.B.; Scott,J.A.; Chin ,S-S. and Amal ,R. 2005. Bactericidal effects of titanium dioxide – based photocatalysts . *Chem. Engineering J.* 113 (1) 55-63.
- Dlgnanl, M.C.and Analssie ,E. 2004.Human fusariosis .*Clin .Microbiol . Infect.*10 : 67-75.
- Doll, T.E. and Frimmel, F.H. 2005 .Cross flow microfiltration with periodical back washing for photocatalytic degradation of pharmaceutical and diagnostic residues – evaluation of the photocatalytic activity of TiO₂ . *Water Research* .39 (5): 847 -854 .
- Fox, M .A. and Dulay, M.T.1993 . Heterogeneous photocatalysis .*Chem.Rev.*93 :341-357.
- Fujishima, A.; Hashimoto, K. and Watanabe, T.1999 .TiO₂ Photocatalysis : fundamentals and application .BKC., Inc. Tokyo, Japan .
- Gassim , F –Al .G.; Alkhateeb, A. N.; Hussein ,F.H. 2004. Photocatalytic oxidation of benzyl alcohol using pure and sensitized anatase. The 8th Arab International World Renewable energy conference and exhibition 8-10 March. King. Bahrain .461-471 .
- Grow, N.R. and Gadd, G.M.1995.The growing fungus .Chapman and Hall ,London. PP.48 .
- Hammel, K.E.; Kapich ,A.N.; Jensen Jr., K.A. and Ryan ,Z.C. 2002. Reactive oxygen species as agents of wood decay by fungi. *Enzyme and Microbial Technol.*30 (4) 445-453 .
- Hansen, H.N. and Smith, R.E.1932 .The mechanisms of variation in imperfect fungi : *Botrytic cinerea*. *Phytopathology*.37 (4) 369-371.
- Higgen, R.A.1973. Engineering metallurgy ,1st ed. Univ. Press LTD .Englaned.P.373.
- Hoffmann, M. R.; Martin ,S. T.; Choi, W. and Bahnemann, D. W. 1995. Environmental applications of semiconductor photocatalysis. *Chem. Rev.* 95 : 69–96.
- Hur, J.S.; Oh, S.O.; Lim, K.M. ; Jung, J.S.; Kim, J.W. and Koh, Y.J. 2005.Novel effects of TiO₂ photocatalytic ozonation on control of post harvest fungal spoilage of kiwifruit post harvest . *Bio.Technol.*35 (1) 109-113.
- Kamat , P.V. 1993. Photochemistry on nonreactive and reactive (semiconductor) surfaces . *Chem. Rev.* 93.267-300 .
- Kim, T. ;Park, K,K.; Jeoung, T. ; Kim, S. and Cho, S. 2006. Sterilization of pathogenic bacteria using titanium dioxide photocatalyst .The Annual Meeting ,San Francisco .CA.
- Kiraly ,Z and Solymosy , F. 1974. Methods in plant pathology . Elsevier Scientific Publishing Comp.New York. P.507.
- Lawton , L.A.; Robertson, P.K.J.; Cornish, B.J.P.A. and Jaspars ,M.1999. Detoxification of microcystins (Cyanobacterial hepatoxins).using TiO₂ photocatalytic oxidation. *Environ . Sci.Technol.*33 :771-775.
- Lee, S-H. 2004. Photocatalytic nano - composites based on TiO₂ and carbon nanotubs. Ph. D.Thesis .Univ. Florida .P.94.
- Leonard, J. M. and Blackford, V. L. 1949. Fungus – Inhibitive proprieties of bromo acetamides. *J. Bacteriol.* 57 :339 - 347.
- Lonnen, J.; Kilvington, S.; Kehoe, S.C.; Al-touati; F .and Mcguigan ,K.G. 2005. Solar and photocatlytic disinfection of protozoan , fungal and bacterial microbes in drinking water .*Water Research* .39 (5) 877-883 .
- Makowski, A. and Wardas, W. 2001.Photocatalytic degradation of toxins secreted to water by cyanobacteria and unicellular algae and photocatlytic degradation of the cells of selected microorganism . *Current Topics in Biophysics* 25 (1) ,19 – 25 .
- Maneerat, C. and Hayata, Y. 2006. Antifungal Activity of TiO₂ Photocatalysis Against *Penicillium expansum* in Fruit testes . *International J. Food Microbiol.*107 (2) 99-103.
- Maness,P.; Smolinski , S.; Blake, D.M.; Huang, Z.; Wolfrum , E.J.and Jacoby ,W.A. 1999. Bactericidal activity of photocatalytic TiO₂ reaction : toward an understanding of its killing mechanism .*Appl. Environ. Microbiol.* 65 (9) : 4094 - 4098 .
- Matsunaga, T.;Tomada, R.; Nakajima,T.and Wake,H.1985.Photochemical sterilization of microbial cells by semiconductor powders . *FEMS Microbiol. Lett.* 29 (1-2) : 211-214.

- Matsunaga , T., Tomada, R.; Nakajima,Y. ; Nakmura ,N. and Komine, T. 1988. Continuous - sterilization system hat uses photosemiconductor powders .Appl. Environ. Microbiol. 54 (6) 1330- 1333 .
- Neiland ,J. B.,1982 .Microbial envelope protein to iron .Ann. Rev. Microbial. 36 .285 .
- Ohmori,K. and Gottlieb,D.1965.Development respiratory enzyme activities during spore germination .Phytopathology 55:1328-1336.
- Robertson , P.K.J. ; Lawton ,L.A. ;Munch ,B. and Rhouzade J. 1997.Processes influencing the destruction of microorganism –LR by TiO₂ photocatalysis . J. Chem. Soc. Commun . 4.393 - 394 .
- Saito, T .; Iwase , T.; Horie, J and Morioka, T.1992 .Mode of photocatalytic bactericidal action of powdered semiconductor TiO₂ on mutants streptococci .J. Photochem. Photobio. B. 14 (4) : 369 -379 .
- Sichel ,C. ; DeCara , M.; Tello , J .; Blance , J.;Fernandez – Ibanez , P.2007.Solar photocatalytic disinfection of agricultured pathogenic fungi : *Fusarium* species Appl.. Catal. B: Environ. 74 (1-2) 152 – 160 .
- Singh, V. P.; Singh, H. B. and Singh, R. B. 1980. The fungicidal effect of neem (*Azadirachta indica*).
- Steel, R.G.D. and Torrie, J.H.1960. Principles and procedures of S-statistics .McGraw-Hill Book Comp.,Inc.London.p.481.
- Tachikawa, T.; Kawai, K.; Asanoi ,Y.; Tojo ,S.; Sugimoto, A.; Fujitsuka ,M. and Majima,T. 2008. Photocatalytic cleavage of single TiO₂ / DNA Nano conjugates .Chemi . Europ. J. 14 (5) 1492 -1498 .
- Youngman ,R.J .1984 .Oxygen activation : is the hydroxyl radical a lwayses biologically relevant ? Trends Biochem .Sci . 9. 280-283 .
- Wong, Y.W.H ;Yuen,C.W.M.; Leung, M.Y.S. ;Ku, S.K.A. and Lam,H.L.I.2006.Selected app - lication of nanotechnology in textiles .AUTEX Research J. 6 (1)1-8 .